gibberellins), amino acids and peptides (hydrolysis products), carbohydrates, vitamins (including C and Riboflavin-B2), micronutrients, humic and fulvic acids.

RIZOCYN contains plant growth regulators (cytokinins, auxins,

RIZOCYN is essential in all the following cases:

• Promotion of the root in all types of plantlets. • Overcoming the transplanting shock and ensuring the soonest

possible establishment of all transplanting plants in the soil.

 Stimulation of the beneficial microbial activity in the soil which increases the bioavailability of the nutrients in the soil.

 Promotion of the seed germination. • Improvement of the function of the root system in periods of

intense stress caused by various factors such as frost, drought etc. Supplement the organic matter.

Better aeration of the roots.

• Increase of the cell divisions in the root.



1 Ermou & Theotokopoulou str., 144 52 Metamorphosis, Tel. +30 210 284 5891, Fax. +30 210 281 7971 Web Site: www.humofert.gr E-mail: info@humofert.gr

Net Content 250 cc Natural biostimulator for the rapid growth of a voluminous and healthy root system

NATURAL ROOT BIOSTIMULANT

APPLICATION RATES - USE Tomato, Cucumber, Eggplant, Pepper, Watermelon, Melon, Strawberry,

rest of vegetables: Greenhouse crops: 10-20 1/ha. Apply right after transplanting with the first watering. Repeat after 2 weeks. Open field crops: 5-10 l/ha. Apply with the transplanting water or right after transplanting with the first watering. Repeat after 3 weeks. Tobacco: 1 l of the product/250 l of planting water. Apply during

planting with the transplanting water. Potato: Dipping of the potato seed in a solution of a dilution ratio 250 ml/100 l of water. Spray on the leaves at an application rate 250

Vineyards, Ornamental plants, Fruit trees: Dipping of the roots and

cuttings in a solution 250-500 ml/100 l of water for 3 minutes before planting. Alternatively apply 3-10 ml per plant or cutting after transplanting by watering with an appropriate amount of water. Lawn: 150-200 ml/100 square meters. Seedbeds: 150-200 ml/100 l of water.

ml/100 l of water at tuber set.

Nurseries: 1-21/1,0001 of water.

COMPATIBILITY-NOTES

It is compatible with all fertilizers and technical products. A compatibility test is always recommended before mixing in the tank. It is not toxic even if the recommended application rates are exceeded by

SHAKE WELL BEFORE ANY USE